

NON-COMMUTATIVE CHERN CHARACTERS OF COMPACT QUANTUM GROUPS

DO NGOC DIEP, ADEREMI O. KUKU, AND NGUYEN QUOC THO

ABSTRACT. In this paper we compute the K-theory (algebraic and topological) and entire periodic cyclic homology for compact quantum groups, define Chern characters between them and show that the Chern characters in both topological and algebraic cases are isomorphisms modulo torsion.

INTRODUCTION

Non commutative differential geometry appeared as some quantization of classical differential geometry. In particular, different cyclic (co)homologies are non-commutative versions of the classical de Rham (co)homology of (differential forms or) currents. Quantum groups were introduced, at first as Hopf bialgebra envelop of Lie groups and later as special cases of quasi-triangular Drinfeld algebras or monoidal categories. The theory is a subject of intensive current research.

Many results have been obtained for quantum groups as analogues of the corresponding classical results for Lie groups, especially the theory of highest weight representations. For compact quantum groups, the representation theory is characterized by the highest weight representations, just as in the case of representation theory of compact Lie groups. For the compact Lie groups, the K -groups, de Rham cohomology groups and the Chern characters between them have been well studied (see [13], [19]-[20]). More recently the Chern characters from K-theory to entire cyclic homology of such groups have also been studied (see [9]). However such studies have not yet been done for quantum groups. The aim of this paper is to compute the non-commutative Chern characters for compact quantum groups in a C^* -algebra setting.

Algebraic and topological K -theories of Banach algebras have been well studied and known to have such nice properties as homotopy invariance, Morita

Date: July 17, 1998.

1991 Mathematics Subject Classification. Primary 22E45; Secondary 19, 43, 58.

Key words and phrases. Quantum group, group C^* -algebras, cyclic homology, K-theory, de Rham currents.

The first author was supported in part by the International Centre for Theoretical Physics, the IMU Commission on Development and Exchange, and Vietnam National Foundation for Research in Fundamental Sciences.

invariance, excision, etc... At the same time, the entire cyclic cohomology $\mathrm{HE}^*(A)$, defined by A. Connes with the non-commutative Chern characters ch_{C^*} , given by the pairing $K_*(A) \times \mathrm{HE}^*(A) \rightarrow \mathbf{C}$ has been rather difficult to compute for various concrete algebras and indeed the authors are not aware of any computations involving algebras other than $A = \mathbf{C}$.

In this paper we define an entire cyclic homology $\mathrm{HE}_*(A)$ by considering a direct system of ideals of A with ad -invariant trace and using the pairing of the entire cochains of A. Connes at the level of ideals with elements of $K_*(A)$

We now the structure of the paper. In Section 1, we at first recall definitions and main properties of compact quantum groups and their representations. More precisely, if G is a complex algebraic group with compact real form, $\mathfrak{g}$ its Lie algebra, ε a positive real number, we define a C^* -algebraic compact quantum group $C_\varepsilon^*(G)$ as the C^* -completion of the $*$ -algebra $\mathcal{F}_\varepsilon(G)$ with respect to the C^* -norm, where $\mathcal{F}_\varepsilon(G)$ is the quantized Hopf algebra of functions which is the Hopf subalgebra of the Hopf algebra dual to the quantized universal enveloping algebra $U(\mathfrak{g})$, generated by matrix elements of the $U(\mathfrak{g})$ modules of type $\mathbf{1}$ (see [6]). We proved that

$$C_\varepsilon^*(G) \cong \mathbf{C}(\mathbf{T}) \oplus \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \mathcal{K}(\mathbf{H}_{w,t}) dt$$

where $\mathcal{K}(\mathbf{H}_{w,t})$ is the elementary algebra of compact operators in separable infinite-dimensional Hilbert space $\mathbf{H}_{w,t}$ and W is the Weyl group of G with respect to a maximal torus $\mathbf{T}$.

In Section 2, we provide the definition and main properties of entire periodic cyclic homology of non-commutative involutive Banach algebras. We prove in particular that when the C^* -algebra A can be presented as direct limit $\varinjlim I_\alpha$, where I_α are ideals of A admitting an ad -invariant trace then $\mathrm{HE}_*(A) = \varinjlim \mathrm{HE}_*(I_\alpha)$.

Section 3 is devoted to Chern characters. We first compute the K -groups of $C_\varepsilon^*(G)$, the $\mathrm{HE}_*(C_\varepsilon^*(G))$. Therefore we define the Chern characters from $K_*(C_\varepsilon^*(G))$ to $\mathrm{HE}_*(C_\varepsilon^*(G))$ for any involutive Banach algebra A endowed with a family of ideals with ad -invariant trace. When $A = C_\varepsilon^*(G)$ we prove that

$$\mathrm{HE}_*(C_\varepsilon^*(G)) \cong \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}})$$

and

$$K_*(C_\varepsilon^*(G)) \cong K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong K^*(\mathcal{N}_{\mathbf{T}}).$$

Finally we show that there is a commutative diagram

$$\begin{array}{ccc}
K_*(C_\varepsilon^*(G)) & \xrightarrow{ch_{C^*}} & \mathrm{HE}_*(C_\varepsilon^*(G)) \\
\cong \downarrow & & \downarrow \cong \\
K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) & \xrightarrow{ch_{CQ}} & \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \\
\cong \downarrow & & \downarrow \cong \\
K^*(\mathcal{N}_{\mathbf{T}}) & \xrightarrow{ch} & H_{DR}^*(\mathcal{N}_{\mathbf{T}}),
\end{array}$$

we deduce that ch_{C^*} is an isomorphism modulo torsion.

In Section 4 we at first identify our definition of $\mathrm{HP}_*(C_\varepsilon^*(G))$ with Cuntz-Quillen definition, and then compute $K_*^{alg}(C_\varepsilon^*(G))$, $\mathrm{HP}(C_\varepsilon^*(G))$ and prove that the Chern character ch_{alg} from $K_{alg}^*(C_\varepsilon^*(G))$ to $\mathrm{HP}_*(C_\varepsilon^*(G))$ is an isomorphism modulo torsion.

Notes on Notation. For any compact space X , we write $K^*(X)$ for the $\mathbf{Z}/2$ graded topological K-theory of X . We use Swan's theorem to identify $K^*(X)$ with $\mathbf{Z}/2$ graded $K_*(\mathbf{C}(X))$. For any involutive Banach algebra A , $K_*(A)$, $\mathrm{HE}_*(A)$, $\mathrm{HP}_*(A)$ are $\mathbf{Z}/2$ graded algebraic or topological K-group of A , entire cyclic homology, and periodic cyclic homology of A , respectively. If $\mathbf{T}$ is a maximal torus of a compact group G , with the corresponding Weyl group W , write $\mathbf{C}(\mathbf{T})$ for the algebra of complex valued functions on $\mathbf{T}$. We use the standard notations from the root theory such as P , P^+ for the positive highest weights, etc.... We denote by $\mathcal{N}_{\mathbf{T}}$ the normalizer of $\mathbf{T}$ in G , by $\mathbf{N}$ the set of natural numbers, $\mathbf{R}$ the field of real numbers and $\mathbf{C}$ the field of complex numbers, $\ell_A^2(\mathbf{N})$ the standard ℓ^2 space of square integrable sequences of elements from A , and finally by $C_\varepsilon^*(G)$ we denote the compact quantum group.

1. COMPACT QUANTUM GROUPS AND THEIR REPRESENTATIONS

In this section we present a summary of some notions and results concerning compact quantum groups and their representations for later use. The main reference is the book [6]

Let G be a complex algebraic group with Lie algebra $\mathfrak{g} = \mathrm{Lie} G$. There is a one-to-one correspondence between:

1. real forms of G ,
2. Hopf *-structures on the universal enveloping algebra $U(\mathfrak{g})$ and
3. Hopf *-structures on the dual $\mathcal{F}(G) := U(\mathfrak{g})'$.

The compact real form K can be characterized as the set of fixed points of conjugate-linear involution of the algebra $\mathcal{F}(G)$ of representative functions of G .

In the Hopf algebra dual $U_\varepsilon(\mathfrak{g})'$, one considers the Hopf sub-algebra $\mathcal{F}_\varepsilon(G)$, generated by the matrix elements of finite-dimensional $U_\varepsilon(\mathfrak{g})$ -module of type **1**, see Definition 13.1.1 from [6].

If ε is real and $\varepsilon \neq -1$, there exists a unique Haar functional on $\mathcal{F}_\varepsilon(G)$ which can be used to prove results similar to those for classical Lie groups, e.g. the Schur orthogonality relations. From this one can effectively study the unitary representations of compact quantum groups, see ([6], Ch. 13).

In the classical theory, there is a (one-dimensional) unitary representation of the algebra $\mathcal{F}(G)$ associated to every point of the real form $G_{\mathbf{R}}$ the corresponding involution. Since in the quantum case there are no “points”, the study of unitary representations replaces the study of the space itself. For compact quantum groups the unitary representations of $\mathcal{F}_\varepsilon(G)$ are parameterized by pairs (w, t) , where w is an element of the Weyl group W of G and t is an element of a fixed maximal torus of the compact real form of G .

Let $\lambda \in P^+$, $V_\varepsilon(\lambda)$ be the irreducible $U_\varepsilon(\mathfrak{g})$ -module of type **1** with the highest weight λ . Then $V_\varepsilon(\lambda)$ admits of a positive definite hermitian form $(\cdot, \cdot)$, such that $(xv_1, v_2) = (v_1x^*v_2)$ for all $v_1, v_2 \in V_\varepsilon(\lambda)$, $x \in U(\mathfrak{g})$.

Let $\{v_\mu^\nu\}$ be an orthogonal basis for weight space $V_\varepsilon(\lambda)_\mu$, $\mu \in P^+$. Then $\cup\{v_\mu^\nu\}$ is an orthogonal basis for $V_\varepsilon(\lambda)$. Let $C_{\nu,s;\mu,r}^\lambda(x) := (xv_\mu^r, v_\nu^s)$ be the associated matrix element of $V_\varepsilon(\lambda)$. Then the matrix elements $C_{\nu,s;\mu,r}^\lambda$ (where λ runs through P^+ , while (μ, r) and (ν, s) runs independently through the index set of a basis of $V_\varepsilon(\lambda)$) form a basis of $\mathcal{F}_\varepsilon(G)$ (see [6]).

Now every irreducible $*$ -representations of $\mathcal{F}_\varepsilon(\mathrm{SL}_2(\mathbf{C}))$ is equivalent to a representation belonging to one of the following two families, each of which is parameterized by $\mathbf{S}^1 = \{t \in \mathbf{C} \mid |t| = 1\}$ – (i) the family of one-dimensional representations τ_t , (ii) the family π_t of representations in $\ell^2(\mathbf{N})$ (see [6], page 437, Proposition 13.1.8).

Moreover, there exists a surjective homomorphism $\mathcal{F}_\varepsilon(G) \rightarrow \mathcal{F}(\mathrm{SL}_2(\mathbf{C}))$ induced by the natural inclusion $\mathrm{SL}_2(\mathbf{C}) \hookrightarrow G$ and by composing the representation π_{-1} of $\mathcal{F}(\mathrm{SL}_2(\mathbf{C}))$ with this homomorphism we obtain a representation of $\mathcal{F}_\varepsilon(G)$ in $\ell^2(\mathbf{N})$ denoted by π_{s_i} where s_i appears in the reduced decomposition $w = s_{i_1}s_{i_2}\dots s_{i_k}$. More precisely, $\pi_{s_i} : \mathcal{F}_\varepsilon(G) \rightarrow \mathcal{L}(\ell^2(\mathbf{N}))$ is of class CCR (see [8]), i.e. its image is dense in the ideal of compact operators in $\mathcal{L}(\ell^2(\mathbf{N}))$.

The representation τ_t is one-dimensional and is of the form

$$\tau_t(C_{\nu,s;\mu,r}^\lambda) = \delta_{r,s}\delta_{\mu\nu} \exp(2\pi\sqrt{-1}\mu(x)),$$

if

$$t = \exp(2\pi\sqrt{-1}x) \in \mathbf{T}, \text{ for } x \in \mathfrak{t} = \mathrm{Lie} \mathbf{T},$$

see §13.1.D from [6].

The unitary representations of compact quantum groups are classified as follows.

- Every irreducible unitary representation of $\mathcal{F}_\varepsilon(G)$ on a Hilbert space is a completion of a unitarizable highest weight representation. Moreover, two such representations are equivalent if and only if they have the same weight, ([6], Theorem 13.1.7).
- The highest weight representations can be described as follows. Let $w = s_{i_1} s_{i_2} \dots s_{i_k}$ be a reduced decomposition of an element w of the Weyl group W . Then, (i) the Hilbert space tensor product

$$\rho_{w,t} = \pi_{s_{i_1}} \otimes \dots \otimes \pi_{s_{i_k}} \otimes \tau_t$$

is an irreducible $*$ -representation of $\mathcal{F}_\varepsilon(G)$ which is associated to the Schubert cell $\mathbf{S}_w$; (ii) up to equivalence, the representation $\rho_{w,t}$ does not depend on the choice of the reduced decomposition of w ; (iii) every irreducible $*$ -representation of $\mathcal{F}_\varepsilon(G)$ is equivalent to some $\rho_{w,t}$, see ([6], Theorem 13.1.9).

Moreover, one can show that

$$\bigcap_{(w,t) \in W \times \mathbf{T}} \ker \rho_{w,t} = \{e\},$$

i.e. the representation

$$\bigoplus_{w \in W} \int_{\mathbf{T}}^{\oplus} \rho_{w,t} dt$$

is a faithful representation and

$$\dim \rho_{w,t} = \begin{cases} 1, & \text{if } w = e, \\ \infty, & \text{if } w \neq e \end{cases}$$

(see Remarks [1] after Theorem 13.1.9 from [6].)

We recall now the definition of C^* -algebraic compact quantum group. Let G be a complex algebraic group with compact real form.

Definition 1.1 ([6], Definition. 13.3.1). The C^* -algebraic compact quantum group $C_\varepsilon^*(G)$ is the C^* -completion of the $*$ -algebra $\mathcal{F}_\varepsilon(G)$ with respect to the C^* -norm

$$\|f\| := \sup_{\rho} \|\rho(f)\| \quad (f \in \mathcal{F}_\varepsilon(G)),$$

where ρ runs through the $*$ -representations of $\mathcal{F}_\varepsilon(G)$ and the norm on the right-hand side is the operator norm.

We have the following result about the structure of compact quantum groups.

Theorem 1.2.

$$C_\varepsilon^*(G) \cong \mathbf{C}(\mathbf{T}) \oplus \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \mathcal{K}(\mathbf{H}_{w,t}) dt,$$

where $\mathbf{C}(\mathbf{T})$ is the algebra of continuous function on $\mathbf{T}$ and $\mathcal{K}(\mathbf{H})$ is the ideal of compact operators in a separable Hilbert space $\mathbf{H}$

Proof. Let $w = s_{i_1} \cdot s_{i_2} \cdot \dots \cdot s_{i_k}$ be a reduced decomposition of the element $w \in W$ into a product of reflections. By construction, the representations $\rho_{w,t} = \pi_{s_{i_1}} \otimes \dots \otimes \pi_{s_{i_k}} \otimes \tau_t$ where π_{s_i} is the composition of the homomorphism of $\mathcal{F}_\varepsilon(G)$ onto $\mathcal{F}_\varepsilon(\mathrm{SL}_2(\mathbf{C}))$ and the representation π_{-1} of $\mathcal{F}_\varepsilon(\mathrm{SL}_2(\mathbf{C}))$ in the Hilbert space $\ell^2(\mathbf{N})$. Thus we have

$$\pi_{s_i} : C_\varepsilon^*(G) \longrightarrow C_\varepsilon^*(\mathrm{SL}_2(\mathbf{C})) \xrightarrow{\pi_{-1}} \mathcal{L}(\ell^2(\mathbf{N}))$$

The last one is CCR (see Dixmier [8]) and we have

$$\pi_{s_i}(C_\varepsilon^*(G)) \cong \mathcal{K}(\mathbf{H}_{s_i}), \quad \tau_t(C_\varepsilon^*(G)) \cong \mathbf{C},$$

$$\rho_{w,t}(C_\varepsilon^*(G)) \cong \mathcal{K}(\mathbf{H}_{s_1}) \otimes \dots \otimes \mathcal{K}(\mathbf{H}_{s_k}) \otimes \mathbf{C} \cong \mathcal{K}(\mathbf{H}_{w,t}),$$

where $\mathbf{H}_{w,t} = \mathbf{H}_{s_1} \otimes \dots \otimes \mathbf{H}_{s_k} \otimes \mathbf{C}$. (Here the tensor product of Banach algebras always means the tensor with the minimal norm.) $\square$

2. STABILITY OF ENTIRE HOMOLOGY OF NON-COMMUTATIVE DE RHAM CURRENTS

We briefly recall in this subsection a definition of operator K-functors. Following G. G. Kasparov, a Fredholm representation of a C^* -algebra A is a triple (π_1, π_2, F) , consisting of $*$ -representations $\pi_1, \pi_2 : A \rightarrow \mathcal{L}(\mathcal{H}_B)$ and a Fredholm operator $F \in \mathcal{F}(\mathcal{H}_B)$, admitting an adjoint operator, on the Hilbert C^* -module $\mathcal{H}_B = \ell_B^2$ over C^* -algebra B , satisfying the relations

$$\pi_1(a)F - F\pi_2(a) \in \mathcal{K}_B,$$

where $\mathcal{K}_B$ is the ideal of compact (adjoint-able) C^* -module endomorphisms of $\mathcal{H}_B$. The classes of homotopy invariance and unitary equivalence of Fredholm modules form the so called Kasparov operator K-group $KK^*(A, B)$. Herewith put $A = C_\varepsilon^*(G)$, $B = \mathbf{C}$, where G is a compact Lie group. In this case, we have $K_*(C_\varepsilon^*(G)) = KK^*(C_\varepsilon^*(G), \mathbf{C})$, where $K_*(A)$ is algebraic K -group of A . From now on we shall use $K_*(A)$ to denote either topological or algebraic K -theory of the Banach algebra A . Note that

$$KK_0(A, \mathbf{C}) \cong K_0(A),$$

$$KK_1(A, \mathbf{C}) \cong K_1(A).$$

Let us recall now the definition and properties of entire homology of Banach algebras. We refer the reader to [10] for a more detailed exposition.

Let A be an involutive Banach algebra. Recall that A. Connes defined entire cyclic cohomology $\mathrm{HE}^*(A)$ and a pairing

$$K_*(A) \times \mathrm{HE}^*(A) \rightarrow \mathbf{C}.$$

Also M. Khalkhali [14] proved Morita and homotopy invariance of $\mathrm{HE}^*(A)$. We now define the entire homology $\mathrm{HE}_*(A)$ as follows: Given a collection $\{I_\alpha\}_{\alpha \in \Gamma}$ of ideals in A , equipped with a so called ad_A -invariant trace

$$\tau_\alpha : I_\alpha \rightarrow \mathbf{C},$$

satisfying the properties:

1. τ_α is a *continuous linear* functional, normalized as $\|\tau_\alpha\| = 1$,
2. τ_α is *positive* in the sense that

$$\tau_\alpha(a^*a) \geq 0, \forall \alpha \in \Gamma,$$

where the map $a \mapsto a^*$ is the involution defining the involutive Banach algebra structure, i.e. an anti-hermitian endomorphism such that $a^{**} = a$

3. τ_α is *strictly positive* in the sense that $\tau_\alpha(a^*a) = 0$ iff $a = 0$, for every $\alpha \in \Gamma$.
4. τ_α is *ad_A -invariant* in the sense that

$$\tau_\alpha(xa) = \tau_\alpha(ax), \forall x \in A, a \in I_\alpha,$$

then we have for every $\alpha \in \Gamma$ a scalar product

$$\langle a, b \rangle_\alpha := \tau_\alpha(a^*b)$$

and also an inverse system $\{I_\alpha, \tau_\alpha\}_{\alpha \in \Gamma}$. Let $\bar{I}_\alpha$ be the completion of I_α with respect to the scalar product, defined above and $\widetilde{\bar{I}_\alpha}$ denote $\bar{I}_\alpha$ with formally adjoined unity element. Define $C^n(\widetilde{\bar{I}_\alpha})$ as the set of $n+1$ -linear maps $\varphi : (\widetilde{\bar{I}_\alpha})^{\otimes(n+1)} \rightarrow \mathbf{C}$. There exists a Hilbert structure on $(\widetilde{\bar{I}_\alpha})^{\otimes(n+1)}$ and we can identify $C_n((\widetilde{\bar{I}_\alpha})) := \mathrm{Hom}(C^n(\widetilde{\bar{I}_\alpha}), \mathbf{C})$ with $C^n(\widetilde{\bar{I}_\alpha})$ via an anti-isomorphism.

For $I_\alpha \subseteq I_\beta$, we have a well-defined map

$$D_\alpha^\beta : C^n(\widetilde{\bar{I}_\alpha}) \rightarrow C^n(\widetilde{\bar{I}_\beta}),$$

which makes $\{C^n(\widetilde{\bar{I}_\alpha})\}$ into a direct system. Write $Q = \varinjlim C^n(\widetilde{\bar{I}_\alpha})$ and observe that it admits a Hilbert space structure, see [10]-[11]. Let $C_n(A) := \mathrm{Hom}(\varinjlim C^n(\widetilde{\bar{I}_\alpha}), \mathbf{C}) = \mathrm{Hom}(\varinjlim C_n(\widetilde{\bar{I}_\alpha}), \mathbf{C})$ which is anti-isomorphic to $\varinjlim_\alpha C^n(\widetilde{\bar{I}_\alpha})$. So we have finally

$$C_n(A) = \varinjlim C_n(\widetilde{\bar{I}_\alpha}).$$

Let

$$b, b' : C^n(\widetilde{\bar{I}_\alpha}) \rightarrow C^{n+1}(\widetilde{\bar{I}_\alpha}),$$

$$N : C^n(\widetilde{\bar{I}_\alpha}) \rightarrow C^n(\widetilde{\bar{I}_\alpha}),$$

$$\lambda : C^n(\widetilde{\bar{I}_\alpha}) \rightarrow C^n(\widetilde{\bar{I}_\alpha}),$$

$$S : C^{n+1}(\widetilde{\bar{I}_\alpha}) \rightarrow C^n(\widetilde{\bar{I}_\alpha})$$

be defined as in A. Connes [3]. We adopt the notations in [14]. Denote by $b^*, (b')^*, N^*, \lambda^*, S^*$ the corresponding adjoint operators. Note also that for each I_α we have the same formulae for adjoint operators for homology as Connes obtained for cohomology.

We now have a bi-complex

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \downarrow (-b')^* & & \downarrow b^* & & \downarrow (-b')^* \\ \mathcal{C}(A) : & \dots & \xleftarrow{1-\lambda^*} \varinjlim_\alpha C_1(\tilde{I}_\alpha) & \xleftarrow{N^*} \varinjlim_\alpha C_1(\tilde{I}_\alpha) & \xleftarrow{1-\lambda^*} \varinjlim_\alpha C_1(\tilde{I}_\alpha) & \xleftarrow{N^*} \dots \\ & & \downarrow (-b')^* & & \downarrow b^* & & \downarrow (-b')^* \\ & \dots & \xleftarrow{1-\lambda^*} \varinjlim_\alpha C_0(\tilde{I}_\alpha) & \xleftarrow{N^*} \varinjlim_\alpha C_0(\tilde{I}_\alpha) & \xleftarrow{1-\lambda^*} \varinjlim_\alpha C_0(\tilde{I}_\alpha) & \xleftarrow{N^*} \dots \end{array}$$

with $d_v := b^*$ in the even columns and $d_v := (-b')^*$ in the odd columns, $d_h := 1 - \lambda^*$ from odd to even columns and $d_h := N^*$ from even to odd columns, where $*$ means the corresponding adjoint operator. Now we have

$$\text{Tot}(\mathcal{C}(A))^{\text{even}} = \text{Tot}(\mathcal{C}(A))^{\text{odd}} := \oplus_{n \geq 0} C_n(A),$$

which is periodic with period two. Hence, we have

$$\oplus_{n \geq 0} C_n(A) \begin{array}{c} \xleftarrow{\partial} \\ \xrightarrow{\partial} \end{array} \oplus_{n \geq 0} C_n(A)$$

where $\partial = d_v + d_h$ is the total differential.

Definition 2.1. Let $\text{HP}_*(A)$ be the homology of the total complex $(\text{Tot}\mathcal{C}(A))$. It is called the *periodic cyclic homology* of A .

Note that this $\text{HP}_*(A)$ is, in general different from the $\text{HP}_*(A)$ of Cuntz-Quillen, because we used the direct limit of periodic cyclic homology of ideals. But in special cases, when the whole algebra A is one of these ideals with ad_A -invariant trace, (e.g. the commutative algebras of complex-valued functions on compact spaces), we return to the Cuntz-Quillen HP_* , which we shall use later.

Definition 2.2. An even (or odd) chain $(f_n)_{n \geq 0}$ in $\mathcal{C}(A)$ is called *entire* if the radius of convergence of the power series $\sum_n \frac{n!}{[\frac{n}{2}]!} \|f_n\| z^n$, $z \in \mathbf{C}$ is infinite.

Let $C^e(A)$ be the sub-complex of $C(A)$ consisting of entire chains. Then we have a periodic complex.

Theorem 2.3. *Let*

$$\text{Tot}(C^e(A))^{\text{even}} = \text{Tot}(C_e(A))^{\text{odd}} := \bigoplus_{n \geq 0} C_n^e(A),$$

where $C_n^e(A)$ is the entire n -chain. Then we have a complex of entire chains with the total differential $\partial := d_v + d_h$

$$\begin{array}{ccc} & \partial & \\ & \longleftarrow & \\ \bigoplus_{n \geq 0} C_n^e(A) & & \bigoplus_{n \geq 0} C_n^e(A) \\ & \longrightarrow & \\ & \partial & \end{array}$$

Definition 2.4. The homology of this complex is called also the *entire homology* and denoted by $\text{HE}_*(A)$.

Note that this entire homology is defined through the inductive limits of ideals with ad-invariance trace.

In [10]-[11], the main properties of this theory, namely

- Homotopy invariance,
- Morita invariance and
- Excision,

were proved and hence that HE_* is a generalized homology theory.

Lemma 2.5. *If the Banach algebra A can be presented as a direct limit $\varinjlim_{\alpha} I_{\alpha}$ of a system of ideals I_{α} with ad-invariant trace τ_{α} , then*

$$K_*(A) = \varinjlim_{\alpha} K_*(I_{\alpha}),$$

$$\text{HE}_*(A) = \varinjlim_{\alpha} \text{HE}_*(I_{\alpha}).$$

Proof. Let us consider $K_*(A) = K_0(A) \oplus K_1(A)$. By definition

$$K_0(A) := \{e^* = e = e^2 \in \text{Mat}_{\infty}(A)\},$$

where

$$\begin{aligned} \text{Mat}_{\infty}(A) &= \varinjlim_n \text{Mat}_n(A) = \varinjlim_n \text{Mat}_n(\varinjlim_{\alpha} I_{\alpha}) = \varinjlim_n \varinjlim_{\alpha} \text{Mat}_n(I_{\alpha}) = \\ &= \varinjlim_{\alpha} \varinjlim_n \text{Mat}_n(I_{\alpha}) = \varinjlim_{\alpha} \text{Mat}_{\infty}(I_{\alpha}). \end{aligned}$$

This means that $K_0(A) = \varinjlim_{\alpha} (I_{\alpha})$. By the same argument we can prove for the K_1 groups:

$$\begin{aligned} K_1(A) &= \text{GL}_{\infty}(A) / (\text{GL}_{\infty}(A), \text{GL}_{\infty}(A)) \\ &\cong \text{GL}_{\infty}(\varinjlim_{\alpha} I_{\alpha}) / (\text{GL}_{\infty}(\varinjlim_{\alpha} I_{\alpha}), \text{GL}_{\infty}(\varinjlim_{\alpha} I_{\alpha})) \\ &\cong \varinjlim_{\alpha} \text{GL}(I_{\alpha}) / (\text{GL}(I_{\alpha}), \text{GL}(I_{\alpha})) \\ &\cong \varinjlim_{\alpha} K_*(I_{\alpha}). \end{aligned}$$

We consider now the $\mathcal{C}(\tilde{I}_\alpha)$ -bi-complex

$$\mathcal{C}(\tilde{I}_\alpha) : \begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \downarrow (-b')^* & & \downarrow b^* & & \downarrow (-b')^* \\ \dots & \xleftarrow{1-\lambda^*} & C_1(\tilde{I}_\alpha) & \xleftarrow{N^*} & C_1(\tilde{I}_\alpha) & \xleftarrow{1-\lambda^*} & C_1(\tilde{I}_\alpha) \xleftarrow{N^*} \dots \\ & & \downarrow (-b')^* & & \downarrow b^* & & \downarrow (-b')^* \\ \dots & \xleftarrow{1-\lambda^*} & C_0(\tilde{I}_\alpha) & \xleftarrow{N^*} & C_0(\tilde{I}_\alpha) & \xleftarrow{1-\lambda^*} & C_0(\tilde{I}_\alpha) \xleftarrow{N^*} \dots \end{array}$$

The connecting maps of the direct system

$$\tilde{I}_\alpha \longrightarrow \tilde{I}_\beta \longrightarrow \tilde{I}_\gamma, \text{ for } \alpha \leq \beta \leq \gamma$$

induces the same connecting maps for complexes

$$\mathcal{C}(\tilde{I}_\alpha) \longrightarrow \mathcal{C}(\tilde{I}_\beta) \longrightarrow \mathcal{C}(\tilde{I}_\gamma), \text{ for } \alpha \leq \beta \leq \gamma$$

and finally we have for the corresponding total complexes:

$$\varinjlim_\alpha \text{Tot}\mathcal{C}(\tilde{I}_\alpha) = \bigoplus_\alpha \text{Tot}\mathcal{C}(\tilde{I}_\alpha) / \sim.$$

It is easy to check the following claims:

$$\mathcal{C}(\bigoplus_\alpha \tilde{I}_\alpha) \cong \bigoplus_\alpha \mathcal{C}(\tilde{I}_\alpha),$$

$$\text{Tot}\mathcal{C}(\bigoplus_\alpha \tilde{I}_\alpha) \cong \bigoplus_\alpha \text{Tot}\mathcal{C}(\tilde{I}_\alpha),$$

and finally

$$\text{Tot}\mathcal{C}(\varinjlim_\alpha \tilde{I}_\alpha) \cong \varinjlim_\alpha \text{Tot}\mathcal{C}(\tilde{I}_\alpha).$$

From the last claim, it is easy to see that a cycle of the complex $\text{Tot}\mathcal{C}(\varinjlim_\alpha \tilde{I}_\alpha)$ can be presented as a sequence of cycles as an element from $\varinjlim_\alpha \text{Tot}\mathcal{C}(\tilde{I}_\alpha)$. The lemma is therefore proved. $\square$

The following result from K-theory is well-known:

Theorem 2.6. *The entire homology of non-commutative de Rham currents admits the following stability property*

$$K_*(\mathcal{K}(\mathbf{H})) \cong K_*(\mathbf{C}),$$

$$K_*(A \otimes \mathcal{K}(\mathbf{H})) \cong K_*(A),$$

where $\mathbf{H}$ is a separable Hilbert space and A is an arbitrary Banach space.

The similar result is true for entire homology HE_* :

Theorem 2.7. *The entire homology of non-commutative de Rham currents admits the following stability property*

$$\mathrm{HE}_*(\mathcal{K}(\mathbf{H})) \cong \mathrm{HE}_*(\mathbf{C}),$$

$$\mathrm{HE}_*(A \otimes \mathcal{K}(\mathbf{H})) \cong \mathrm{HE}_*(A),$$

where $\mathbf{H}$ is a separable Hilbert space and A is an arbitrary Banach space.

Proof. It is easy to see that $\mathcal{K}(\mathbf{H})$ is a direct limit of matrix algebras $\mathrm{Mat}_n(\mathbf{C})$ as

$$\mathcal{K}(\mathbf{H}) = \varinjlim_n \mathrm{Mat}_n(\mathbf{C}),$$

and thus

$$A \otimes \mathcal{K}(\mathbf{H}) = A \otimes \varinjlim_n \mathrm{Mat}_n(\mathbf{C}) = \varinjlim_n A \otimes \mathrm{Mat}_n(\mathbf{C}).$$

The theorem is therefore deduced from the previous lemma. $\square$

3. NON-COMMUTATIVE CHERN CHARACTERS

Let A be a involutive Banach algebra. We construct a non-commutative character

$$ch_{C^*} : K_*(A) \rightarrow \mathrm{HE}_*(A)$$

and later show that when $A = C_\varepsilon^*(G)$, this Chern character reduces up to isomorphism to classical Chern character on the normalizers of maximal compact tori.

Let A be an involutive Banach algebra with unity.

Theorem 3.1. *There exists a Chern character*

$$ch_{C^*} : K_*(A) \rightarrow \mathrm{HE}_*(A).$$

Proof. We first recall that there exists a pairing

$$K_n(A) \times C^n(\widetilde{I}_\alpha) \rightarrow \mathbf{C}$$

due to A. Connes, see [4]. Hence there exists a map $K_n(A) \xrightarrow{C_n} \mathrm{Hom}(C^n(A), \mathbf{C})$. So, by 1.1, we have for each α , a map $K_n(A) \xrightarrow{C_n^\alpha} \mathrm{Hom}(C^n(\widetilde{I}_\alpha), \mathbf{C})$ and hence a map $K_n(A) \xrightarrow{C_n} \mathrm{Hom}(\varinjlim_\alpha C^n(\widetilde{I}_\alpha), \mathbf{C})$. We now show that C_n induces the Chern map

$$ch : K_n(A) \rightarrow \mathrm{HE}_n(A)$$

Now let e be an idempotent in $M_k(A)$ for some $k \in \mathbf{N}$. It suffices to show that for n even, if $\varphi = \partial\psi$, where $\varphi \in C^n(\widetilde{I}_\alpha)$ and $\psi \in C^{n+1}(\widetilde{I}_\alpha)$, then

$$\langle e, \varphi \rangle = \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \varphi(e, e, \dots, e) = 0.$$

However, this follows from Connes' results in ([4], Lemma 7).

The proof of the case for n odd would also follow from [4]. $\square$

Our next result computes the Chern characters for $A = C_\varepsilon^*(G)$ by reducing it to the classical case.

Theorem 3.2. *Let $\mathbf{T}$ be a fixed maximal torus of G with Weyl group $W := \mathcal{N}_{\mathbf{T}}/\mathbf{T}$. Then the Chern character*

$$ch_{C^*} : K_*(C_\varepsilon^*(G)) \rightarrow \mathrm{HE}_*(C_\varepsilon^*(G))$$

is an isomorphism modulo torsion, i.e.

$$ch_{C^*} : K_*(C_\varepsilon^*(G)) \otimes \mathbf{C} \xrightarrow{\cong} \mathrm{HE}_*(C_\varepsilon^*(G)),$$

which can be identified with the classical Chern character

$$ch : K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \longrightarrow \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$$

that is also an isomorphism modulo torsion, i.e.

$$ch : K_*(\mathcal{N}_{\mathbf{T}}) \otimes \mathbf{C} \xrightarrow{\cong} H_{DR}^*(\mathcal{N}_{\mathbf{T}}).$$

To prove this theorem we first observe:

Lemma 3.3.

$$K_*(C_\varepsilon^*(G)) \cong K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong K^*(\mathcal{N}_{\mathbf{T}}).$$

Proof. We have

$$\begin{aligned} K_*(C_\varepsilon^*(G)) &= K_*(\mathbf{C}(\mathbf{T}) \oplus \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^\oplus \mathcal{K}(\mathbf{H}_{w,t}) dt) \\ &= K_*(\mathbf{C}(\mathbf{T}) \oplus K_*(\bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^\oplus \mathcal{K}(\mathbf{H}_{w,t}) dt)) \\ &= K_*(\mathbf{C}(\mathbf{T})) \oplus K_*(\bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^\oplus \varinjlim_n \mathrm{Mat}_n(\mathbf{C}) dt) \\ &= K_*(\mathbf{C}(\mathbf{T})) \oplus K_*(\varinjlim_n \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^\oplus \mathrm{Mat}_n(\mathbf{C}) dt) \\ &= K_*(\mathbf{C}(\mathbf{T})) \oplus \varinjlim_n K_*(\mathbf{C}((W - \{e\}) \times \mathbf{T}) \otimes \mathrm{Mat}_n(\mathbf{C})) \\ &\cong K_*(\mathbf{C}(\mathbf{T})) \oplus K_*(\mathbf{C}((W - \{e\}) \times \mathbf{T})) \\ &= K_*(\mathbf{C}(\mathbf{T}) \oplus \mathbf{C}((W - \{e\}) \times \mathbf{T})) \\ &= K_*(\mathbf{C}(W \times \mathbf{T})) \\ &= K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}/\mathbf{T} \times \mathbf{T})) \\ &= K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})), \end{aligned}$$

by using Morita invariance property. The last one is isomorphic with the topological K-theory $K^*(\mathcal{N}_{\mathbf{T}})$. $\square$

Next we have

Lemma 3.4.

$$\mathrm{HE}_*(C_\varepsilon^*(G)) \cong \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{DR}^*(\mathcal{N}_{\mathbf{T}}).$$

Proof. The proof is analogous to that of 3.3. So we have

$$\begin{aligned}
\mathrm{HE}_*(C_\varepsilon^*(G)) &= \mathrm{HE}_*(\mathbf{C}(\mathbf{T}) \oplus \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \mathcal{K}(\mathbf{H}_{w,t}) dt) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathbf{T}) \oplus \mathrm{HE}_*(\bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \mathcal{K}(\mathbf{H}_{w,t}) dt)) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathbf{T})) \oplus \mathrm{HE}_*(\bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \varinjlim_n \mathrm{Mat}_n(\mathbf{C}) dt) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathbf{T})) \oplus \mathrm{HE}_*(\varinjlim_n \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^{\oplus} \mathrm{Mat}_n(\mathbf{C}) dt) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathbf{T})) \oplus \varinjlim_n \mathrm{HE}_*(\mathbf{C}((W - \{e\}) \times \mathbf{T}) \otimes \mathrm{Mat}_n(\mathbf{C})) \\
&\cong \mathrm{HE}_*(\mathbf{C}(\mathbf{T})) \oplus \mathrm{HE}_*(\mathbf{C}((W - \{e\}) \times \mathbf{T})) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathbf{T}) \oplus \mathbf{C}((W - \{e\}) \times \mathbf{T})) \\
&= \mathrm{HE}_*(\mathbf{C}(W \times \mathbf{T})) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}/\mathbf{T} \times \mathbf{T})) \\
&= \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})),
\end{aligned}$$

by using Morita invariance property of HE_* . Now, since $\mathbf{C}(\mathcal{N}_{\mathbf{T}})$ is a commutative $\mathbf{C}$ -algebra, then by Cuntz-Quillen's result [7], we have a canonical isomorphism $\mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}})$, since $\mathbf{C}(\mathcal{N}_{\mathbf{T}})$ is smooth. Moreover by a result of Khalkhali [14]-[15] $\mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$; hence the result. $\square$

Proof of Theorem 3.2

Now consider the commutative diagram

$$\begin{array}{ccc}
K_*(C_\varepsilon^*(G)) & \xrightarrow{ch_{C^*}} & \mathrm{HE}_*(C_\varepsilon^*(G)) \\
\alpha \downarrow & & \downarrow \gamma \\
(I) \quad K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) & \xrightarrow{ch_{CQ}} & \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \\
\beta \downarrow & & \downarrow \delta \\
K^*(\mathcal{N}_{\mathbf{T}}) & \xrightarrow{ch} & H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}})
\end{array}$$

Now, $K_*(C_\varepsilon^*(G)) \cong K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$ by Lemma 3.3. Also $\mathrm{HE}_*(C_\varepsilon^*(G)) \cong \mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$ by Lemma 3.4. By a result of Khalkhali [14]-[15], we have $\mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong \mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$. Moreover $\mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}})$, since the algebra $\mathbf{C}(\mathcal{N}_{\mathbf{T}})$ is smooth in the sense of Cuntz-Quillen (see [7]). Hence, $\mathrm{HE}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}})$. Furthermore, $K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong K^*(\mathcal{N}_{\mathbf{T}})$ by the well-known Swan' theorem. So all the vertical maps are isomorphisms. It is well known that the bottom map ch is also an isomorphism modulo torsion, i.e.

$$ch : K^*(\mathcal{N}_{\mathbf{T}}) \otimes \mathbf{C} \xrightarrow{\cong} H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}}).$$

Hence ch_{CQ} and ch_{C^*} are isomorphisms modulo torsion. The theorem is therefore proved.

4. ALGEBRAIC VERSION

Let $C_\varepsilon^*(G)$ be a compact quantum group, $\mathrm{HP}_*(C_\varepsilon^*(G))$ the periodic cyclic homology introduced in §2. Since $C_\varepsilon^*(G) = \mathbf{C}(\mathbf{T}) \oplus \bigoplus_{e \neq w \in W} \int_{\mathbf{T}}^\oplus \mathcal{K}(\mathbf{H}_{w,t}) dt$

$\mathrm{HP}_*(C_\varepsilon^*(G))$ coincides with the $\mathrm{HP}_*(C_\varepsilon^*(G))$ defined by J. Cuntz-D. Quillen [7].

J. Cuntz and D. Quillen [7] defined the so called X -complexes of $\mathbf{C}$ -algebras and then used some ideas of Fedosov product to define algebraic Chern characters. We now briefly recall their definitions. For a (non-commutative) associative $\mathbf{C}$ -algebra A , consider the space of even non-commutative differential forms $\Omega^+(A) \cong RA$, equipped with the Fedosov product

$$\omega_1 \circ \omega_2 := \omega_1 \omega_2 - (-1)^{|\omega_1|} d\omega_1 d\omega_2,$$

see [7]. Consider also the ideal $IA := \bigoplus_{k \geq 1} \Omega^{2k}(A)$. It is easy to see that $RA/IA \cong A$ and that RA admits the universal property that any based linear map $\rho : A \rightarrow M$ can be uniquely extended to a derivation $D : RA \rightarrow M$. The derivations $D : RA \rightarrow M$ bijectively correspond to lifting homomorphisms from RA to the semi-direct product $RA \oplus M$, which also bijectively correspond to linear map $\bar{\rho} : \bar{A} = A/\mathbf{C} \rightarrow M$ given by

$$a \in \bar{A} \mapsto D(\rho a).$$

From the universal property of $\Omega^1(RA)$, we obtain a bi-module isomorphism

$$RA \otimes \bar{A} \otimes RA \cong \Omega^1(RA).$$

As in [7], let $\Omega^- A = \bigoplus_{k \geq 0} \Omega^{2k+1} A$. Then we have

$$\Omega^- A \cong RA \otimes \bar{A} \cong \Omega^1(RA)_\# := \Omega^1(RA)/[(\Omega^1(RA), RA)].$$

J. Cuntz and D. Quillen proved

Theorem 4.1. ([7], Theorem 1): *There exists an isomorphism of $\mathbf{Z}/(2)$ -graded complexes*

$$\Phi : \Omega A = \Omega^+ A \oplus \Omega^- A \cong RA \oplus \Omega^1(RA)_\#,$$

such that

$$\Phi : \Omega^+ A \cong RA,$$

is defined by

$$\Phi(a_0 da_1 \dots da_{2n} = \rho(a_1) \omega(a_1, a_2) \dots \omega(a_{2n-1}, a_{2n}),$$

and

$$\Phi : \Omega^- A \cong \Omega^1(RA)_\#,$$

$$\Phi(a_0 da_1 \dots da_{2n+1}) = \rho(a_1) \omega(a_1, a_2) \dots \omega(a_{2n-1}, a_{2n}) \delta(a_{2n+1}).$$

With respect to this identification, the product in RA is just the Fedosov product on even differential forms and the differentials on the X -complex

$$X(RA) : \quad RA \cong \Omega^+ A \rightarrow \Omega^1(RA)_\# \cong \Omega^- A \rightarrow RA$$

become the operators

$$\beta = b - (1 + \kappa)d : \Omega^- A \rightarrow \Omega^+ A,$$

$$\delta = -N_{\kappa^2}b + B : \Omega^+ A \rightarrow \Omega^- A,$$

where $N_{\kappa^2} = \sum_{j=0}^{n-1} \kappa^{2j}$, $\kappa(da_1 \dots da_n) := da_n \dots da_1$.

Let us denote by $IA \triangleleft RA$ the ideal of even non-commutative differential forms of order ≥ 2 . By the universal property of Ω^1

$$\Omega^1(RA/IA) = \Omega^1 RA / ((IA)\Omega^1 RA + \Omega^1 RA.(IA) + dIA).$$

Since $\Omega^1 RA = (RA)dRA = dRA.(RA)$, then $\Omega^1 RA(IA) \cong IA\Omega^1 RA \pmod{[RA, \Omega^1 R]}$.

$$\Omega^1(RA/IA)_{\#} = \Omega^1 RA / ([RA, \Omega^1 RA] + IA.dRA + dIA).$$

For IA -adic tower $RA/(IA)^{n+1}$, we have the complex

$$\mathcal{X}(RA/(IA)^{n+1}) : \quad RA/IA^{n+1} \leftarrow \Omega^1 RA / ([RA, \Omega^1 RA] + (IA)^{n+1}dRA + d(IA)^{n+1}).$$

Define

$$\begin{aligned} \mathcal{X}^{2n+1}(RA, IA) : \quad RA/(IA)^{n+1} &\rightarrow \Omega^1 RA / ([RA, \Omega^1 RA] + (IA)^{n+1}dRA + d(IA)^{n+1}) \\ &\rightarrow RA/(IA)^{n+1}, \end{aligned}$$

$$\begin{aligned} \mathcal{X}^{2n}(RA, IA) : \quad RA/((IA)^{n+1} + [RA, IA^n]) &\rightarrow \Omega^1 RA / ([RA, \Omega^1 RA] + d(IA)^n dRA) \\ &\rightarrow RA/((IA)^{n+1} + [RA, IA^n]). \end{aligned}$$

One has

$$\begin{aligned} b((IA)^n dIA) &= [(IA)^n, IA] \subset (IA)^{n+1}, \\ d(IA)^{n+1} &\subset \sum_{j=0}^n (IA)^j d(IA)(IA)^{n-j} \subset (IA)^n dIA + [RA, \Omega^1 RA]. \end{aligned}$$

and hence

$$\begin{aligned} \mathcal{X}^1(RA, IA) &= X(RA, IA), \\ \mathcal{X}^0(RA, IA) &= (RA/IA)_{\#}. \end{aligned}$$

There is a sequence of maps between complexes

$$\dots \rightarrow X(RA/IA) \rightarrow \mathcal{X}^{2n+1}(RA, IA) \rightarrow \mathcal{X}^{2n}(RA, IA) \rightarrow X(RA/IA) \rightarrow \dots$$

We have the inverse limits

$$\hat{X}(RA, IA) := \varprojlim X(RA/(IA)^{n+1}) = \varprojlim \mathcal{X}^n(RA, IA).$$

Remark that

$$\begin{aligned} \mathcal{X}^q &= \Omega A / F^q \Omega A, \\ \hat{X}(RA/IA) &= \hat{\Omega} A. \end{aligned}$$

We quote the second main result of J. Cuntz and D. Quillen ([7], Thm2), namely:

$$H_i \hat{\mathcal{X}}(RA, IA) = \text{HP}_i(A).$$

We now apply this machinery to our case. First we have the following algebraic analogy of the Lemma for HP_* .

Lemma 4.2.

$$\text{HP}_*(C_\varepsilon^*(G)) \cong \text{HP}_*(C(\mathcal{N}_T)) \cong H_{\text{DR}}^*(\mathcal{N}_T).$$

Proof. By analogy with the proof of the previous proposition, we also have

$$\begin{aligned} \text{HP}_*(C_\varepsilon^*(G)) &= \text{HP}_*(C(T) \oplus \bigoplus_{e \neq w \in W} \int_T^\oplus \mathcal{K}(\mathbf{H}_{w,t}) dt) \\ &= \text{HP}_*(C(T) \oplus \text{HP}_*(\bigoplus_{e \neq w \in W} \int_T^\oplus \mathcal{K}(\mathbf{H}_{w,t}) dt)) \\ &= \text{HP}_*(C(T)) \oplus \text{HP}_*(\bigoplus_{e \neq w \in W} \int_T^\oplus \varinjlim_n \text{Mat}_n(C) dt) \\ &= \text{HP}_*(C(T)) \oplus \text{HP}_*(\varinjlim_n \bigoplus_{e \neq w \in W} \int_T^\oplus \text{Mat}_n(C) dt) \\ &= \text{HP}_*(C(T)) \oplus \varinjlim_n \text{HP}_*(C((W - \{e\}) \times T) \otimes \text{Mat}_n(C)) \\ &\cong \text{HP}_*(C(T)) \oplus \text{HP}_*(C((W - \{e\}) \times T)) \\ &= \text{HP}_*(C(T) \oplus C((W - \{e\}) \times T)) \\ &= \text{HP}_*(C(W \times T)) \\ &= \text{HP}_*(C(\mathcal{N}_T/T \times T)) \\ &= \text{HP}_*(C(\mathcal{N}_T)), \end{aligned}$$

by using Morita invariance property of HP_* . The last one is following Cuntz-Quillen for the commutative $\mathbf{C}$ -algebra A , we have a canonical isomorphism from periodic cyclic homology to the $\mathbf{Z}/(2)$ -graded de Rham homology, when A is smooth. $\square$

Now, for each idempotent $e \in M_n(A)$ there is an unique element $x \in M_n(\widehat{RA})$. Then the element

$$\tilde{e} := x + (x - \frac{1}{2}) \sum_{n \geq 1} \frac{2^n(2n-1)!!}{n!} (x - x^{2n})^{2n} \in M_n(\widehat{RA})$$

is a lifting of e to an idempotent matrix in $M_n(\widehat{RA})$. Then the map $[e] \mapsto \text{tr}(\tilde{e})$ defines the map $K_0(A) \rightarrow H_0(X(\widehat{RA})) = \text{HP}_0(A)$. To an element $g \in \text{GL}_n(A)$ one associates an element $p \in \text{GL}(\widehat{RA})$ and to the element g^{-1} an element $q \in \text{GL}_n(\widehat{RA})$ then put

$$x = 1 - qp, \text{ and } y = 1 - pq.$$

And finally, to each class $[g] \in \text{GL}_n(A)$ one associates

$$\text{tr}(g^{-1}dg) = \text{tr}(1-x)^{-1}d(1-x) = d(\text{tr}(\log(1-x))) = -\text{tr} \sum_{n=0}^{\infty} x^n dx \in \Omega^1(A)_\#.$$

Then $[g] \rightarrow \text{tr}(g^{-1}dg)$ defines the map $K_1(A) \rightarrow HH_1(A) = H_1(X(\widehat{RA})) = \text{HP}_1(A)$.

Definition 4.3. Let $\mathrm{HP}^*(I_n)$ be the periodic cyclic cohomology defined by Cuntz-Quillen. Then the pairing

$$K_*^{\mathrm{alg}}(C^*(G)) \times \bigcup_n \mathrm{HP}^*(I_n) \rightarrow \mathbf{C},$$

where $I_n = \mathbf{C}(\mathbf{T}) \oplus \bigoplus_{\mathbf{T}}^{\oplus} \mathrm{Mat}_n(\mathbf{C}) \cong \mathbf{C}(\mathbf{T}) \oplus \mathbf{C}((W - \{e\}) \times \mathbf{T}) \otimes \mathrm{Mat}_n(\mathbf{C})$, defines an algebraic non-commutative Chern character

$$ch_{\mathrm{alg}} : K_*^{\mathrm{alg}}(C_{\varepsilon}^*(G)) \rightarrow \mathrm{HP}_*(C_{\varepsilon}^*(G)),$$

which gives us a variant of non-commutative Chern characters with values in HP_* -groups.

We close this section with an algebraic analog of theorem 3.2.

Theorem 4.4. *Let G be a compact Lie group and $\mathbf{T}$ a fixed maximal compact torus of G . Then in the notations of 4.3, the Chern character*

$$ch_{\mathrm{alg}} : K_*(C_{\varepsilon}^*(G)) \rightarrow \mathrm{HP}_*(C_{\varepsilon}^*(G))$$

is an isomorphism modulo torsion, i.e.

$$ch_{\mathrm{alg}} : K_*(C_{\varepsilon}^*(G)) \otimes \mathbf{C} \rightarrow \mathrm{HP}_*(C_{\varepsilon}^*(G))$$

which can be identified with the classical Chern,

$$ch : K^*(\mathcal{N}_{\mathbf{T}}) \longrightarrow H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}}),$$

which is equivalent to the Cuntz-Quillen character

$$ch_{CQ} : K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \rightarrow \mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}}))$$

which are also isomorphisms modulo torsion, i.e.

$$ch : K^*(\mathcal{N}_{\mathbf{T}}) \otimes \mathbf{C} \xrightarrow{\cong} H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}}),$$

$$ch_{CQ} : K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \otimes \mathbf{C} \xrightarrow{\cong} \mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})).$$

Proof. First note that

$$K_*(C_{\varepsilon}^*(G)) \cong K_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong K^*(\mathcal{N}_{\mathbf{T}}).$$

Next we have

$$\mathrm{HP}_*(C_{\varepsilon}^*(G)) \cong \mathrm{HP}_*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H_{\mathrm{DR}}^*(\mathcal{N}_{\mathbf{T}}).$$

Furthermore, by a result of Cuntz-Quillen for the commutative $\mathbf{C}$ -algebra A , we have a canonical isomorphism from periodic cyclic homology to the $\mathbf{Z}/(2)$ -graded de Rham cohomology, when A is smooth. Hence

$$K^*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})) \cong H^*(\mathbf{C}(\mathcal{N}_{\mathbf{T}})).$$

Now we have a commutative diagram

$$\begin{array}{ccc}
K_*(C_\varepsilon^*(G)) & \xrightarrow{ch_{alg}} & HP_*(C_\varepsilon^*(G)) \\
\cong \downarrow & & \downarrow \cong \\
K_*(C(\mathcal{N}_T)) & \xrightarrow{ch_{CQ}} & HP_*(C(\mathcal{N}_T)) \\
\cong \downarrow & & \downarrow \cong \\
K^*(\mathcal{N}_T) & \xrightarrow{ch} & H_{DR}^*(\mathcal{N}_T)
\end{array}$$

where the vertical homomorphisms all are isomorphisms. Hence, ch_{alg} is an isomorphism modulo torsion. $\square$

ACKNOWLEDGMENTS

This work was completed during the stay of the first author as a visiting mathematician at the International Centre for Theoretical Physics, Trieste, Italy. He would like to thank ICTP for the hospitality, without which this work would not have been possible and the IMU Commission of Development and Exchange for a generous grant.

This work is supported in part by the International Centre for Theoretical Physics, Trieste, Italy, the IMU Commission on Development and Exchange and the Vietnam National Foundation for Research in Fundamental Sciences.

REFERENCES

- [1] N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac operators*, Grundlehren der mathematischen Wissenschaften, Corrected 2nd Printing 1996, No. 298, Springer-Verlag, 1992.
- [2] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Annals of Math., **57**(1953), 115–207.
- [3] A. Connes, *Non commutative Geometry*, Academic Press, 1994.
- [4] A. Connes, *Entire cyclic cohomology of Banach algebras and characters of θ -summable Fredholm modules*, K-theory, **1**(1988), 519–548.
- [5] J. Cuntz, *A survey of some aspects of non-commutative geometry*, Mathematisches Inst. Uni. Heidelberg, **35**(1992), pp.1–29.
- [6] V. Chari and A. Pressley, *A guide to Quantum Groups*, Cambridge Univ. Press, 1995.
- [7] J. Cuntz and D. Quillen, *The X complex of the universal extensions*, Preprint Math. Inst. Uni. Heidelberg, 1993.
- [8] J. Dixmier, *C*-algebras*, North-Holland, Amsterdam, 1982.
- [9] D. N. Diep, A. O. Kuku and N. Q. Tho, *Noncommutative Chern characters of compact Lie group C*-algebras*, K-Theory (to appear).
- [10] D. N. Diep and N. V. Thu, *Homotopy invariance of entire current cyclic homology*, Vietnam J. of Math., **25**(1997), No 3, 211–228.

- [11] D. N. Diep and N. V. Thu, *Entire homology of non-commutative de Rham currents*, ICTP, IC/96/214, 1996, 23pp; to appear in Publication of CFCA, Hanoi-City Vietnam National University, 1997.
- [12] F. Feng and B. Tsygan, *Hochschild and cyclic homology of quantum groups*, Commun. Math. Phys. **140**(1991), 481–521.
- [13] L. Hodgkin, *On the K-theory of Lie groups*, Topology, **6**(1967), 1-36.
- [14] M. Khalkhali, *On the entire cyclic cohomology of Banach algebras: I. Morita invariance*, Mathematisches Inst. Uni. Heidelberg, **54**(1992), pp 24.
- [15] M. Khalkhali, *On the entire cyclic cohomology of Banach algebras: II. Homotopy invariance*, Mathematisches Inst. Uni. Heidelberg, **55**(1992), pp 18.
- [16] A. A. Kirillov, *Elements of the Theory of representations*, Springer-Verlag, Berlin-Heidelberg-New York, 1975.
- [17] N. V. Thu, *Morita invariance of entire current cyclic homology*, Vietnam J. Math. (to appear).
- [18] N. V. Thu, *Exactness of entire current cyclic homology*, Acta Math. Vietnamica (to appear).
- [19] T. Watanabe, *On the Chern characters of symmetric spaces related to $SU(n)$* , J. Math. Kyoto Univ. **34**(1994), 149–169.
- [20] T. Watanabe, *Chern characters on compact Lie groups of low rank*, Osaka J. Math. **22**(1985), 463–488.

INSTITUTE OF MATHEMATICS, NATIONAL CENTRE FOR NATURAL SCIENCE AND TECHNOLOGY, P.O. BOX 631, BO HO, 10.000, HANOI, VIETNAM

Current address: International Centre for Theoretical Physics, ICTP P. O. Box 586, 34100, Trieste, Italy

E-mail address: dndiep@ioit.ncst.ac.vn

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS, ICTP P. O. BOX 586, 34100, TRIESTE, ITALY

E-mail address: kuku@ictp.trieste.it

DEPARTMENT OF MATHEMATICS, VINH UNIVERSITY, VINH CITY, VIETNAM

Current address: c/o Institute of Mathematics, National Centre for Natural Science and Technology, P.O. Box 631, Bo Ho, 10.000, Hanoi, Vietnam

E-mail address: dndiep@ioit.ncst.ac.vn